FLIP antibody

Cat. No. GTX26144

Host	Rabbit	
Clonality	Polyclonal	
lsotype	IgG	
Application	WB, ICC/IF, ELISA	
Reactivity	Human, Mouse, Rat	

Reference (1) Package 100 μg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1µg/ml
ICC/IF	10µg/ml
ELISA	Assay dependent
Not tacted in other applications	

Not tested in other applications.

Calculated MW

55 kDa. (<u>Note</u>)

PROPERTIES		
Form	Liquid	
Buffer	PBS	
Preservative	0.02% Sodium azide	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)	
Immunogen	Peptide corresponding to aa 447- 464 at the C-terminus of human FLIP α /FLIPL (accession no. AAC51622)	
Purification	Purified by affinity chromatography	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	




For full product information, images and publications, please visit our <u>website</u>.




Datasheet

DATA IMAGES



GTX26144 ICC/IF Image

ICC/IF analysis of Jurkat cells using GTX26144 FLIP antibody. Dilution : 10 $\mu\text{g/ml}$



GTX26144 ICC/IF Image

ICC/IF analysis of K562 cells using GTX26144 FLIP antibody. Red : Primary antibody Blue : DAPI Dilution : 5 μg/ml



GTX26144 ICC/IF Image

Immunocytochemical staining of FLIPa in Jurkat cells with FLIPa antibody (GTX26144) at 10µg/ml.



GTX26144 WB Image

Western blot analysis of FLIP in HeLa (H), Jurkat (J), and K562 (K) whole cell lysate with FLIP antibody (GTX26144) at 1ug/ml. Positive control: Whole cell lysate from HeLa cells.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 03 Page 2 of 2