

Goat Anti-Rat IgG (Fc fragment) antibody, F(ab')₂ fragment, pre-adsorbed (PE)**Cat. No. GTX26259**

Host	Goat
Clonality	Polyclonal
Isotype	IgG F(ab') ₂
Applications	ICC/IF, FCM
Reactivity	Rat

Package
500 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:100-1:250
FCM	1:100-1:250

Not tested in other applications.

Product Note Pre-adsorbed with Bovine, Horse and Human serum proteins. May react with immunoglobulins from other species.

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Rat IgG Fc fragment
Purification	Purified by antigen-affinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, pepsin digestion and chromatographic separation. From serum
Conjugation	Phycoerythrin (PE) Wavelength
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).