

# Donkey Anti-Goat IgG antibody, Fab fragment (TxRd)

**Cat. No. GTX26523**

<b>Host</b>	Donkey
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG Fab
<b>Applications</b>	ICC/IF, FCM, Dot, ELISA
<b>Reactivity</b>	Goat

**Package**  
1 mg

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:1000-1:5000
FCM	1:500-1:2500
Dot	Assay dependent
ELISA	1:10000-1:50000

Not tested in other applications.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
<b>Preservative</b>	0.01% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Goat IgG whole molecule
<b>Purification</b>	IgG fraction This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, papain digestion and chromatographic separation.
<b>Conjugation</b>	Texas Red (TxRd) <a href="#">Wavelength</a> Ratio : 1.61 molecules TxRd per Donkey IgG Fab molecule.



For full product information, images and publications, please visit our [website](#).

**Note**

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).