

# p27 Kip1 antibody

**Cat. No. GTX26547**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, IP, ELISA
<b>Reactivity</b>	Human

**Package**  
100 µl

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:4000
IP	1:100
ELISA	1:10000-1:40000

Not tested in other applications.

**Calculated MW** 22 kDa. ( [Note](#) )

**Product Note** No reactivity is observed with p27 from mouse.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	20mM Potassium Phosphate, 150mM NaCl
<b>Preservative</b>	0.01% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	80 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Full length recombinant human p27 protein.
<b>Conjugation</b>	Unconjugated


**Note** For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


## DATA IMAGES



### GTX26547 WB Image

WB analysis of MCF-7 whole cell lysate using GTX26547 p27 Kip1 antibody.

Dilution : 1:500




### GTX26547 WB Image

WB analysis of non-transfected and transfected 293T whole cell lysate using GTX26547 p27 Kip1 antibody.

Loading : 30 µg

Dilution : 1:2000



### GTX26547 WB Image

Western blot using GeneTex affinity purified anti-p27 antibody (GTX26547) shows detection of p27 protein in MCF7 whole cell lysate (lanes 1-3) (arrowhead). Separation was achieved using a 4-20% gradient gel.

Blocking occurred using 5% BLOTTO. Primary antibody was diluted 1:500 in 1% BLOTTO. The membrane was washed and reacted with a 1:10,000 dilution of Dylight™ 800 conjugated Goat anti Rabbit IgG.

Molecular weight estimation was made by comparison to prestained MW markers indicated at the left (lane M). Other detection systems will yield similar results.



For full product information, images and publications, please visit our [website](https://www.genetex.com).