

Biotin antibody (FITC)

Cat. No. GTX26650

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	ICC/IF, FCM, Dot
Reactivity	Species independent

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:500-1:2500
FCM	Assay dependent
Dot	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Biotin conjugated to -KLH
Purification	Purified by antigen-affinity chromatography. From serum
Conjugation	Fluorescein isothiocyanate (FITC) Wavelength Ratio : 4.6 molecules FITC per Goat IgG molecule.

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).