

Fluorescein antibody (TxRd)

Cat. No. GTX26654

| | |
|--------------|---------------------|
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, ELISA |
| Reactivity | Species independent |

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | >1:10000 |
| ICC/IF | 1:500-1:2500 |
| ELISA | >1:20000 |

Not tested in other applications.

Properties

| | |
|---------------|--|
| Form | Liquid |
| Buffer | 20mM Potassium Phosphate, 150mM NaCl, 1% BSA |
| Preservative | 0.01% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Fluorescein conjugated to Goat IgG |
| Purification | Purified by antigen-affinity chromatography. From serum |
| Conjugation | Texas Red (TxRd) Wavelength Ratio : 2.9 molecules Texas Red per Goat IgG molecule. |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 06 Page 1 of 1