

Goat Anti-Human IgG antibody, pre-adsorbed (FITC)

Cat. No. GTX27149

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Application	ICC/IF, FACS, ELISA
Reactivity	Human

Package $100 \, \mu g$

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:1000-1:5000
FACS	1:500-1:2500
ELISA	1:10000-1:50000
Notation of the other conflictions	

Not tested in other applications.

Pre-adsorbed with Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat and Sheep serum proteins. May **Product Note** react with immunoglobulins from other species.

PROPERTIES	
Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human IgG whole molecule
Purification	Purified by antigen-affinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities. From serum
Conjugation	Fluorescein isothiocyanate (FITC) Ratio: 2.19 molecules FITC per Goat IgG molecule.



For full product information, images and publications, please visit our website.

Date 2024 / 05 / 13 Page 1 of 2

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com



For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 13 Page 2 of 2