

beta Tubulin antibody [DM1B]

Cat. No. GTX27287

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF
Reactivity	Human, Mouse, Rat, Gerbil

References (1)

Package

100 µg

Applications

Application Note

For IF: Use at a concentration of 4 - 6 µg/ m l. For WB: Use at a concentration of 4 -6µg/ m l. Detects a band of approximately 60 kDa. Optimal dilutions/concentrations should be determined by the researcher.

Properties

Form	Liquid
Buffer	50mM Sodium Phosphate buffer, 0.2% Gelatin
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Native chicken brain microtubules
Purification	Purified IgG
Conjugation	Unconjugated

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).