

CD31 antibody [WM59] (Biotin)

Cat. No. GTX27385

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Applications	FCM
Reactivity	Human, Cynomolgus monkey, Rhesus Monkey

References (28)
Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Recommended dilution Neat-1/10 ted working dilution to label 10 ⁶ cells or 100μl whole blood. s antibody recognizes the human CD31 antigen. Clone WM59 has been mapped to the Ig-like domain 2 (Fawcett et al. 95).
ted working dilution to label 10 ⁶ cells or 100μl whole blood. s antibody recognizes the human CD31 antigen. Clone WM59 has been mapped to the Ig-like domain 2 (Fawcett et al.
s antibody recognizes the human CD31 antigen. Clone WM59 has been mapped to the Ig-like domain 2 (Fawcett et al.
uid
5, 1% BSA
9% Sodium azide
re as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. Fo g-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
mg/ml (Please refer to the vial label for the specific concentration.)
tein A purified m tissue culture supernatant
tin
laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal on nan consumption.
)



For full product information, images and publications, please visit our <u>website</u>.

determine the structure or sequence of the product.

Date 2025 / 12 / 12 Page 1 of 1

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to