

# Parathyroid Hormone antibody

**Cat. No. GTX27843**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, ICC/IF, IHC-P, IHC-Fr, IP
<b>Reactivity</b>	Human

**Package**  
500 µl

## Applications

### Application Note

For WB: Use at a dilution of 1:100-1:200. For IP: Use at an assay dependent dilution. For IHC: Use at a dilution of 1:40-1:80. This antibody can be used on frozen tissue sections and on formalin-fixed, paraffin-embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of 1:40-1:80 with Auto/Probe III. Auto/Zyme (Cat# GTX73183) predigestion or high temperature antigenic unmasking (see instructions below) is required with this antibody prior to application of this antibody with formalin-fixed, paraffin-embedded tissue. The optimum dilution factor should be determined by individual laboratories.

<b>Calculated MW</b>	13 kDa. ( <a href="#">Note</a> )
<b>Product Note</b>	This antibody reacts with parathyroid hormone. Specificity tested by immunodiffusion & immunoelectrophoresis.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 0.2% BSA
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Human intact Parathyroid Hormone
<b>Purification</b>	IgG fraction From polyclonal serum
<b>Conjugation</b>	Unconjugated

### Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).