

CAD antibody

Cat. No. GTX28406

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF
Reactivity	Human, Mouse, Rat

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1µg/ml
ICC/IF	5µg/ml

Not tested in other applications.

Calculated MW 39 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide corresponding to aa 147-164 near the center of murine CAD (accession no. NP_031885). The sequence differs from human DFF40 by two amino acids.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX28406 WB Image

Western blot analysis of DFF40/CAD in HeLa (H), K562 (K), Jurkat (J), and Raji (R) whole cell lysate with DFF40/CAD antibody (GTX28406) at 1µg/ml.



GTX28406 ICC/IF Image

Immunocytochemical staining of DFF40 in Jurkat cells with CAD antibody (GTX28406) at 5µg/ml.



For full product information, images and publications, please visit our [website](https://www.genetex.com).