

BMF antibody

Cat. No. GTX29653

| | |
|--------------|--------------------------|
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, IHC-P, ELISA |
| Reactivity | Human |

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 2.5-5µg/ml |
| ICC/IF | Assay dependent |
| IHC-P | Assay dependent |
| ELISA | Assay dependent |

Not tested in other applications.

Calculated MW 21 kDa. ([Note](#))

Product Note This antibody recognizes an epitope at the C-terminus of human Bmf (Bcl-2-Modifying Factor), a novel BH3-only protein.

Properties

| | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 0.02% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | Peptide corresponding to aa 171- 184 of human Bmf (accession no. NP_277038). |
| Purification | Purified by affinity chromatography |
| Conjugation | Unconjugated |




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX29653 ICC/IF Image


Immunofluorescent staining of Bmf in human kidney tissue with Bmf antibody (GTX29653) at 10µg/ml.



GTX29653 WB Image

WB analysis of HepG2 cell lysate using GTX29653 BMF antibody.

Dilution : 2.5µg/ml (Lane A) and 5µg/ml (Lane B)



GTX29653 WB Image

Western blot analysis of Bmf expression in human HepG2 (lane 1) and 293 (lane 2) cell lysates with Anti-Bmf (Cat. No. GTX29653) at 2 mg /mL.



For full product information, images and publications, please visit our [website](https://www.genetex.com).