

GTX300028

iPSC Antibody Panel -Thomson factor

Product Content

Cat No	Product Name	Reactivity	Applications	Package
GTX100622	Oct4 antibody	Human, Mouse	WB, ICC/IF, FCM, IP	25 µl
GTX100863	Nanog antibody [N3C3]	Human, Mouse, Ceratotherium simum	WB, ICC/IF, IHC-P, IHC-Fr, IP, ELISA, Sandwich ELISA	25 µl
GTX121923	Lin28A antibody	Human, Mouse, Pig	WB, ICC/IF, IHC-P, FCM	25 µl
GTX213110-01	Goat Anti-Rabbit IgG antibody (HRP)	Rabbit	WB, IHC-P, ELISA	25 µl
GTX48037	SSEA-4 antibody [MC-813-70]	Human	WB, ICC/IF, IHC-P, FCM, IP, IHC	25 µl
GTX636505	SOX2 antibody [HL1193]	Human, Mouse, Rat, Cat	WB, ICC/IF, IHC-P, IP	25 µl

Note

For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Oct4 antibody

Cat. No. GTX100622

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, FCM, IP
Reactivity	Human, Mouse

References (5)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:20000
ICC/IF	1:100-1:1000
FCM	1:50-1:200
IP	1:500-1:1000

Not tested in other applications.

Calculated MW 39 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.32 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Oct4. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTx100622 IP Image

Oct4 antibody immunoprecipitates Oct4 protein in IP experiments. IP Sample: cell lysate/extract of Oct4 gene transfected 293T cells A. Cell lysate/extract of transfected 293T cell B. Control with 2 µg of preimmune rabbit IgG C. Immunoprecipitation of Oct4 by 2 µg of Oct4 antibody (GTx100622) 12% SDS-PAGE The immunoprecipitated Oct4 protein was detected by Oct4 antibody (GTx100622) diluted at 1:1000. EasyBlot anti-rabbit IgG (GTx221666-01) was used as a secondary reagent.




GTx100622 WB Image

Non-transfected (-) and transfected (+) NT2D1 whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with Oct4 antibody (GTx100622) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



GTx100622 ICC/IF Image

Oct4 antibody detects Oct4 protein at nucleus by immunofluorescent analysis. Sample: NT2D1 cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: Oct4 stained by Oct4 antibody (GTx100622) diluted at 1:500. Red: phalloidin, a cytoskeleton marker, diluted at 1:200. Scale bar= 10 µm.




GTx100622 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed Human ESC, using Oct4(GTx100622) antibody at 1:400 dilution.




For full product information, images and publications, please visit our [website](#).



GTX100622 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed human embryonic stem cell, using Oct4(GTX100622) antibody at 1:100 dilution.



GTX100622 WB Image


Sample (20 µg of whole cell lysate)

A: mouse ESC

12% SDS PAGE

GTX100622 diluted at 1:5000


The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX100622 WB Image

Human tissue extract (30 µg) was separated by 10% SDS-PAGE, and the membrane was blotted with Oct4 antibody (GTX100622) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

The observed M.W. is based on the publication: PMID: 23642364



GTX100622 FCM Image

Flow cytometry on human embryonic stem cells, staining with Oct4 (GTX100622)antibody at 1:50 dilution(green) or rabbit IgG (black).



For full product information, images and publications, please visit our [website](http://www.genetex.com).

**GTX100622 ICC/IF Image**

Immunofluorescence analysis of paraformaldehyde-fixed human embryonic stem cell, using Oct4(GTX100622) antibody at 1:200 dilution.



For full product information, images and publications, please visit our [website](#).

Nanog antibody [N3C3]

Cat. No. GTX100863

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr, IP, ELISA, Sandwich ELISA
Reactivity	Human, Mouse, Ceratotherium simum

References (48)

★★★★★ Review (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	Assay dependent
IHC-Fr	Assay dependent
IP	1:100-1:500
ELISA	1:1000-1:10000
Sandwich ELISA	Assay dependent

Note : Capture: GTX100863 , Detection: GTX89079

Not tested in other applications.

Calculated MW 35 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human NANOG. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 06 / 10 Page 6 of 24

Conjugation Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Note Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX100863 WB Image

Sample (30 µg of whole cell lysate)
 A: HeLa nucleus
 12% SDS PAGE
 GTX100863 diluted at 1:3000
 The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX100863 IP Image

Immunoprecipitation of NANOG protein from NT2D1 whole cell extracts using 5 µg of NANOG antibody [N3C3] (GTX100863).
 Western blot analysis was performed using NANOG antibody [N3C3] (GTX100863) diluted at 1:500.
 EasyBlot anti-Rabbit IgG (GTX221666-01) was used as a secondary reagent.




GTX100863 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed Human ESC, using NANOG(GTX100863) antibody at 1:400 dilution.




For full product information, images and publications, please visit our [website](#).



GTX100863 ELISA Image

ELISA detection of NANOG using GTX100863 for capture at a concentration of 5 µg/mL and GTX89079 for detection at a concentration of 5 µg/mL.




GTX100863 ICC/IF Image

Nanog antibody [N3C3] detects Nanog protein at nucleus by immunofluorescent analysis.

Sample: NT2D1 cells were fixed in 2% paraformaldehyde at RT for 15 min.

Green: Nanog stained by Nanog antibody [N3C3] (GTX100863) diluted at 1:500.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.



GTX100863 ICC/IF Image


Nanog antibody [N3C3] detects Nanog protein by immunofluorescent analysis.

Sample: PANC-1 cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: Nanog stained by Nanog antibody [N3C3] (GTX100863) diluted at 1:500.


Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.

Blue: Fluoroshield with DAPI (GTX30920).



GTX100863 WB Image

The data was published in the 2021 in Cancer Drug Resist. [PMID: 35582384](https://pubmed.ncbi.nlm.nih.gov/35582384/).




GTX100863 WB Image

The data was published in the 2019 in Cancers (Basel). [PMID: 31861872](https://pubmed.ncbi.nlm.nih.gov/31861872/).




For full product information, images and publications, please visit our [website](https://www.genetex.com).




GTX100863 WB Image

The data was published in the journal J Biomed Sci in 2016. [PMID: 27863490](https://pubmed.ncbi.nlm.nih.gov/27863490/)




GTX100863 WB Image

The data was published in the journal Cell Death Dis in 2015. [PMID: 26136074](https://pubmed.ncbi.nlm.nih.gov/26136074/)



GTX100863 WB Image

The data was published in the journal Cell Death Dis in 2015. [PMID: 26136074](https://pubmed.ncbi.nlm.nih.gov/26136074/)




GTX100863 WB Image

The data was published in the journal Elife in 2017. [PMID: 28537559](https://pubmed.ncbi.nlm.nih.gov/28537559/)




For full product information, images and publications, please visit our [website](https://www.genetex.com).




GTX100863 WB Image

The data was published in the journal Elife in 2017. [PMID: 28537559](https://pubmed.ncbi.nlm.nih.gov/28537559/)



GTX100863 ICC/IF Image

The data was published in the journal Oncotarget in 2020. [PMID: 32256979](https://pubmed.ncbi.nlm.nih.gov/32256979/)



GTX100863 ICC/IF Image

The data was published in the journal Nat Commun in 2018. [PMID: 29973581](https://pubmed.ncbi.nlm.nih.gov/29973581/)



For full product information, images and publications, please visit our [website](https://www.genetex.com).

Lin28A antibody

Cat. No. GTX121923

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, FCM
Reactivity	Human, Mouse, Pig

References (1)

★★★★☆ Review (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:10000
ICC/IF	Assay dependent
IHC-P	1:100-1:1000
FCM	1:50-1:200

Not tested in other applications.

Calculated MW 23 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Lin28A. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note
Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES


**GTX121923 FCM Image**

Flow cytometry on human embryonic stem cells, staining with Lin28A (GTX121923) antibody at 1:100 dilution (purple) or rabbit IgG (black).

**GTX121923 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded human gastric cancer, using Lin28A antibody (GTX121923) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min

**GTX121923 WB Image**

Sample (20 µg of whole cell lysate)

A: mouse ESC


12% SDS PAGE

GTX121923 diluted at 1:3000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](http://www.genetex.com).

**GTX121923 WB Image**

Sample (20 µg of whole cell lysate)

A: hESC

12% SDS PAGE

GTX121923 diluted at 1:5000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](#).

Goat Anti-Rabbit IgG antibody (HRP)

Cat. No. GTX213110-01

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Rabbit

References (638)

Package

1 ml

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:5000-1:20000
IHC-P	1:100-1:1000
ELISA	1:5000-1:20000

Not tested in other applications.

Properties

Form	Liquid
Buffer	0.05M Tris, 0.15M NaCl
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Highly purified whole rabbit IgG
Purification	Purified by antigen-affinity chromatography.
Conjugation	Horseradish peroxidase(HRP)


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX213110-01 ELISA Image

Indirect ELISA analysis performed by coating plate with recombinant SARS-CoV-2 (COVID-19) Spike S1 protein, His tag (active) (GTX135817-pro) (52.15-0.81 nM). Coated protein was probed with SARS-CoV-2 (COVID-19) Spike S1 antibody [HL134] (GTX635671) (1 µg/mL). Goat anti-rabbit IgG antibody (HRP) (GTX213110-01) (1:10000) was used to detect bound primary antibody.




GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in mouse intestine by immunohistochemical analysis. Sample: Paraffin-embedded mouse intestine. WBP11 antibody (GTX118654) diluted at 1:500. The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01) Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various amounts of denatured rabbit IgG protein were separated by 12% SDS-PAGE, and the membrane was blotted with HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) diluted at 1:5000.



GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in rat prostate by immunohistochemical analysis. Sample: Paraffin-embedded rat prostate. WBP11 antibody (GTX118654) diluted at 1:500. The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01). Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various whole cell extracts were separated by 10% SDS-PAGE, and the membrane was blotted with alpha Tubulin antibody (GTX112141) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




For full product information, images and publications, please visit our [website](#).




GTx213110-01 WB Image

The data was published in the 2022 in Cancers (Basel). [PMID: 35267540](https://pubmed.ncbi.nlm.nih.gov/35267540/)




GTx213110-01 WB Image

The data was published in the 2022 in Front Aging Neurosci. [PMID: 35264943](https://pubmed.ncbi.nlm.nih.gov/35264943/)



GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](https://pubmed.ncbi.nlm.nih.gov/28281525/)




GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](https://pubmed.ncbi.nlm.nih.gov/28281525/)




For full product information, images and publications, please visit our [website](https://www.genetex.com).




GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](https://pubmed.ncbi.nlm.nih.gov/28281525/)




GTx213110-01 WB Image

The data was published in the 2016 in PLoS One. [PMID: 27486856](https://pubmed.ncbi.nlm.nih.gov/27486856/)




GTx213110-01 WB Image

The data was published in the journal Front Pharmacol in 2018. [PMID: 30087612](https://pubmed.ncbi.nlm.nih.gov/30087612/)



GTx213110-01 WB Image

The data was published in the journal Cell Commun Signal in 2014. [PMID: 25214434](https://pubmed.ncbi.nlm.nih.gov/25214434/)




GTx213110-01 WB Image

The data was published in the journal PLoS One in 2017. [PMID: 28832634](https://pubmed.ncbi.nlm.nih.gov/28832634/)




For full product information, images and publications, please visit our [website](http://www.genetex.com).




GTX213110-01 WB Image

The data was published in the journal PLoS One in 2017. [PMID: 28945763](https://pubmed.ncbi.nlm.nih.gov/28945763/)




GTX213110-01 WB Image

The data was published in the journal Oncoscience in 2015. [PMID: 26097885](https://pubmed.ncbi.nlm.nih.gov/26097885/)




GTX213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](https://pubmed.ncbi.nlm.nih.gov/31885126/)



GTX213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](https://pubmed.ncbi.nlm.nih.gov/31885126/)



GTX213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](https://pubmed.ncbi.nlm.nih.gov/31885126/)



For full product information, images and publications, please visit our [website](https://www.genetex.com).

SSEA-4 antibody [MC-813-70]

Cat. No. GTX48037

Host	Mouse
Clonality	Monoclonal
Isotype	IgG3
Applications	WB, ICC/IF, IHC-P, FCM, IP, IHC
Reactivity	Human

References (6)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
FCM	Assay dependent
IP	Assay dependent
IHC	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS, 0.05% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human embryonal carcinoma cell line 2102Ep
Purification	Purified by affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

**GTX48037 ICC/IF Image**

ICC/IF analysis of H9 cells using GTX48037 SSEA-4 antibody.

Green : Primary antibody

**GTX48037 FCM Image**

FACS analysis of 2102Ep cells using GTX48037 SSEA-4 antibody.

Red : Primary antibody

Green : Isotype control

Antibody amount : 0.1 ug



For full product information, images and publications, please visit our [website](#).

SOX2 antibody [HL1193]

Cat. No. GTX636505

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human, Mouse, Rat, Cat

 Review (2)

Package
 100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent
IP	Assay dependent

Not tested in other applications.

Observed MW (kDa) 34 kDa.

Product Note This antibody may not be suitable for rat IHC-P assay, as its rat reactivity is based on ICC/IF testing.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of human SOX2.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).


Date 2026 / 06 / 10 Page 21 of 24

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX636505 WB Image


Non-transfected (-) and transfected (+) NT2D1 whole cell extract (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with SOX2 antibody [HL1193] (GTX636505) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX636505 WB Image


Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membranes were blotted with SOX2 antibody [HL1193] (GTX636505) diluted at 1:1000 and competitor's antibody (sc-17320) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

*The competitor is not affiliated with GeneTex and does not endorse this product.



GTX636505 WB Image

Mouse ESC extract (50 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with SOX2 antibody [HL1193] (GTX636505) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX636505 IHC-P Image

SOX2 antibody detects SOX2 protein by immunohistochemical analysis.


Sample: Paraffin-embedded rat tissues.

SOX2 stained by SOX2 antibody (GTX636505) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




For full product information, images and publications, please visit our [website](http://www.genetex.com).




GTX636505 ICC/IF Image

SOX2 antibody [HL1193] detects SOX2 protein at nucleus by immunofluorescent analysis.
 Sample: DIV10 rat E8 primary hippocampal neuron cells were fixed in 4% paraformaldehyde at RT for 15 min.
 Green: SOX2 stained by SOX2 antibody [HL1193] (GTX636505) diluted at 1:500.
 Red: Tau, an axon marker, stained by Tau antibody [GT287] (GTX634809) diluted at 1:500.




GTX636505 WB Image

Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with SOX2 antibody [HL1193] (GTX636505) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX636505 WB Image

Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with SOX2 antibody [HL1193] (GTX636505) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX636505 IHC-P Image

SOX2 antibody [HL1193] detects SOX2 protein by immunohistochemical analysis.
 Sample: Paraffin-embedded human tonsil.
 SOX2 stained by SOX2 antibody [HL1193] (GTX636505) diluted at 1:100.
 Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min




GTX636505 IHC-P Image

SOX2 antibody [HL1193] detects SOX2 protein by immunohistochemical analysis.
 Sample: Paraffin-embedded mouse tissues.
 SOX2 stained by SOX2 antibody [HL1193] (GTX636505) diluted at 1:2000.
 Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).

**GTX636505 IHC-P Image**

SOX2 antibody [HL1193] detects SOX2 protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.

SOX2 stained by SOX2 antibody [HL1193] (GTX636505) diluted at 1:2000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).