

GTX300031

The 9-1-1 Complex Antibody Panel

Product Content

Cat No	Product Name	Reactivity	Applications	Package
GTX100078	Rad9 antibody	Human, Mouse, Rat	WB, ICC/IF, IHC-P	25 µl
GTX101777	FEN1 antibody	Human, Mouse, Rat	WB, ICC/IF, IHC-P, IP, IHC, PLA	25 µl
GTX113745	HUS1 antibody	Human, Mouse, Rat	WB, IHC-P	25 µl
GTX213110-01	Goat Anti-Rabbit IgG antibody (HRP)	Rabbit	WB, IHC-P, ELISA	25 µl
GTX637858	PCNA antibody [HL1965]	Human, Mouse, Rat, Zebrafish, Cat, Dog, Japanese Medaka	WB, IHC-P, IHC-Fr, IHC-Wm	25 µl
GTX70317	Rad1 antibody [4126]	Human	WB	25 µl

Note

For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Rad9 antibody

Cat. No. GTX100078

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

References (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 43 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Rad9. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX100078 WB Image

Rad9 antibody detects Rad9 protein by Western blot analysis.

- A. 30 µg 293T whole cell lysate/extract
 - B. 30 µg A431 whole cell lysate/extract
 - C. 30 µg HeLa whole cell lysate/extract
 - D. 30 µg HepG2 whole cell lysate/extract
- 10 % SDS-PAGE

Rad9 antibody (GTX100078) dilution: 1:1000




GTX100078 IHC-P Image

Rad9 antibody detects RAD9A protein at nucleus on HBL435 xenograft by immunohistochemical analysis.

Sample: Paraffin-embedded HBL435 xenograft.

Rad9 antibody (GTX100078) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min




GTX100078 WB Image

Rad9 antibody detects Rad9 protein by Western blot analysis.

- A. 30 µg PC-12 whole cell lysate/extract
- 10 % SDS-PAGE

Rad9 antibody (GTX100078) dilution: 1:500



GTX100078 WB Image


Rad9 antibody detects Rad9 protein by Western blot analysis.

- A. 30 µg Neuro2A whole cell lysate/extract
 - B. 30 µg GL261 whole cell lysate/extract
 - C. 30 µg C8D30 whole cell lysate/extract
 - D. 30 µg NIH-3T3 whole cell lysate/extract
 - E. 30 µg BCL-1 whole cell lysate/extract
- 10 % SDS-PAGE

Rad9 antibody (GTX100078) dilution: 1:500



For full product information, images and publications, please visit our [website](https://www.genetex.com).


GTX100078 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using Rad9 (GTX100078) antibody at 1:100 dilution.



For full product information, images and publications, please visit our [website](#).

FEN1 antibody

Cat. No. GTX101777

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP, IHC, PLA
Reactivity	Human, Mouse, Rat

References (4)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500
IHC	Assay dependent
PLA	Assay dependent

Not tested in other applications.

Calculated MW 43 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.41 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human FEN1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX101777 WB Image

Whole cell extract (30 µg) was separated by 10% SDS-PAGE, and the membrane was blotted with FEN1 antibody (GTX101777) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX101777 IP Image

FEN1 antibody immunoprecipitates FEN1 protein in IP experiments.

IP samples: Jurkat whole cell extract

A. 30 µg Jurkat whole cell extract


B. Control with 4 µg of preimmune Rabbit IgG

C. Immunoprecipitation of FEN1 protein by 4 µg FEN1 antibody (GTX101777)

10 % SDS-PAGE

The immunoprecipitated FEN1 protein was detected by FEN1 antibody (GTX101777) diluted at 1:500.

[EasyBlot anti-rabbit IgG (GTX221666-01) was used as a secondary reagent]



GTX101777 WB Image

FEN1 antibody detects FEN1 protein by western blot analysis.

A. 30 µg PC-12 whole cell lysate/extract


10% SDS-PAGE

FEN1 antibody (GTX101777) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).



GTx101777 WB Image

FEN1 antibody detects FEN1 protein by western blot analysis.

A. 30 µg NIH-3T3 whole cell lysate/extract

B. 30 µg JC whole cell lysate/extract


C. 30 µg BCL-1 whole cell lysate/extract

D. 30 µg Raw264.7 whole cell lysate/extract

10% SDS-PAGE

FEN1 antibody (GTx101777) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



GTx101777 ICC/IF Image


FEN1 antibody detects FEN1 protein at nucleus by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: FEN1 stained by FEN1 antibody (GTx101777) diluted at 1:500.


Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTx628802) diluted at 1:1000.

Blue: Fluoroshield with DAPI (GTx30920).



GTx101777 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed A549, using FEN1(GTx101777) antibody at 1:200 dilution.



GTx101777 IHC-P Image

FEN1 antibody detects FEN1 protein at nucleus on human breast carcinoma by immunohistochemical analysis.

Sample: Paraffin-embedded human breast carcinoma.

FEN1 antibody (GTx101777) diluted at 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](https://www.genetex.com).

HUS1 antibody

Cat. No. GTX113745

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 32 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.71 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human HUS1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES


GTX113745 IHC-P Image

HUS1 antibody detects HUS1 protein at cytosol and nucleus on rat fore brain by immunohistochemical analysis.

Sample: Paraffin-embedded rat fore brain.


HUS1 antibody (GTX113745) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min


GTX113745 IHC-P Image

Immunohistochemical analysis of paraffin-embedded CA922 xenograft , using HUS1(GTX113745) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX113745 WB Image

Sample (30 ug of whole cell lysate)

A: 293T

12% SDS PAGE

GTX113745 diluted at 1:1000


GTX113745 IHC-P Image

HUS1 antibody detects HUS1 protein at cytosol and nucleus on mouse lymph node by immunohistochemical analysis.


Sample: Paraffin-embedded mouse lymph node.

HUS1 antibody (GTX113745) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](https://www.genetex.com).



GTX113745 WB Image

HUS1 antibody detects HUS1 protein by Western blot analysis.

A. 30 μ g C8D30 whole cell lysate/extract

12 % SDS-PAGE

HUS1 antibody (GTX113745) dilution: 1:1000



For full product information, images and publications, please visit our [website](https://www.genetex.com).

Goat Anti-Rabbit IgG antibody (HRP)

Cat. No. GTX213110-01

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Rabbit

References (654)

Package

1 ml

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:5000-1:20000
IHC-P	1:100-1:1000
ELISA	1:5000-1:20000

Not tested in other applications.

Properties

Form	Liquid
Buffer	0.05M Tris, 0.15M NaCl
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Highly purified whole rabbit IgG
Purification	Purified by antigen-affinity chromatography.
Conjugation	Horseradish peroxidase(HRP)

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX213110-01 ELISA Image

Indirect ELISA analysis performed by coating plate with recombinant SARS-CoV-2 (COVID-19) Spike S1 protein, His tag (active) (GTX135817-pro) (52.15-0.81 nM). Coated protein was probed with SARS-CoV-2 (COVID-19) Spike S1 antibody [HL134] (GTX635671) (1 µg/mL). Goat anti-rabbit IgG antibody (HRP) (GTX213110-01) (1:10000) was used to detect bound primary antibody.




GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in mouse intestine by immunohistochemical analysis.
Sample: Paraffin-embedded mouse intestine.
WBP11 antibody (GTX118654) diluted at 1:500.
The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01)
Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various amounts of denatured rabbit IgG protein were separated by 12% SDS-PAGE, and the membrane was blotted with HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) diluted at 1:5000.



GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in rat prostate by immunohistochemical analysis.
Sample: Paraffin-embedded rat prostate.
WBP11 antibody (GTX118654) diluted at 1:500.
The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01).
Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various whole cell extracts were separated by 10% SDS-PAGE, and the membrane was blotted with alpha Tubulin antibody (GTX112141) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




For full product information, images and publications, please visit our [website](https://www.genetex.com).




GTx213110-01 WB Image

The data was published in the 2022 in Cancers (Basel). [PMID: 35267540](#)




GTx213110-01 WB Image

The data was published in the 2022 in Front Aging Neurosci. [PMID: 35264943](#)



GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](#)




GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](#)




For full product information, images and publications, please visit our [website](#).




GTx213110-01 WB Image

The data was published in the 2017 in Nat Commun. [PMID: 28281525](#)




GTx213110-01 WB Image

The data was published in the 2016 in PLoS One. [PMID: 27486856](#)




GTx213110-01 WB Image

The data was published in the journal Front Pharmacol in 2018. [PMID: 30087612](#)



GTx213110-01 WB Image

The data was published in the journal Cell Commun Signal in 2014. [PMID: 25214434](#)




GTx213110-01 WB Image

The data was published in the journal PLoS One in 2017. [PMID: 28832634](#)




For full product information, images and publications, please visit our [website](#).




GTx213110-01 WB Image

The data was published in the journal PLoS One in 2017. [PMID: 28945763](#)




GTx213110-01 WB Image

The data was published in the journal Oncoscience in 2015. [PMID: 26097885](#)




GTx213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)



GTx213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)



GTx213110-01 WB Image

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)



For full product information, images and publications, please visit our [website](#).

PCNA antibody [HL1965]

Cat. No. GTX637858

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, IHC-P, IHC-Fr, IHC-Wm
Reactivity	Human, Mouse, Rat, Zebrafish, Cat, Dog, Japanese Medaka

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent
IHC-Fr	Assay dependent
IHC-Wm	Assay dependent

Not tested in other applications.

Observed MW (kDa) 36 kDa.

Product Note

Highly recommended for IHC-P in mouse tissues.

This antibody may not be suitable for zebrafish IHC-P / IHC-Fr assays, as its zebrafish reactivity is based on WB / IHC-Wm testing.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of human PCNA
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

GTX637858 IHC-Fr Image


PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Frozen-sectioned mouse testis.

Green: PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:200.

Red: Vimentin antibody [GT812] (GTX629744) diluted at 1:1000.


Blue: Hoechst 33342 staining.


GTX637858 WB Image

Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with

PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody


(GTX213110-01) was used to detect the primary antibody.


GTX637858 WB Image

Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with

PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody

(GTX213110-01) was used to detect the primary antibody.


GTX637858 WB Image


Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with

PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody


(GTX213110-01) was used to detect the primary antibody.




For full product information, images and publications, please visit our [website](https://www.genetex.com).


GTX637858 WB Image

Whole zebrafish extract (30 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


GTX637858 WB Image

Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody. Corresponding RNA expression data for the same cell lines are based on Human Protein Atlas program.


GTX637858 IHC-P Image


PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded rat tissues.

PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:15000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.


GTX637858 IHC-P Image


PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.

PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:400000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.


GTX637858 IHC-P Image

PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.


PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:400000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.



For full product information, images and publications, please visit our [website](https://www.genetex.com).



GTX637858 IHC-P Image

PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse uterus.

PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:400000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTX637858 IHC-P Image


PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded human tissues.

PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:200000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.



GTX637858 IHC-P Image

PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraffin-embedded human tonsil.

PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:200000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX637858 IHC-Wm Image

PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.

Sample: Paraformaldehyde-fixed 2 days-post-fertilization zebrafish embryo.

Green: PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:1000.

Antigen Retrieval: Tris-HCl buffer, pH 9.0, 20 min at 70°C


GTX637858 IHC-Wm Image

PCNA antibody [HL1965] detects PCNA protein by immunohistochemical analysis.


Sample: Paraffin-embedded 2 days-post-fertilization zebrafish embryo.

Green: PCNA stained by PCNA antibody [HL1965] (GTX637858) diluted at 1:1000.

Antigen Retrieval: Tris-HCl buffer, pH 9.0, 20 min at 70°C



For full product information, images and publications, please visit our [website](https://www.genetex.com).



GTX637858 WB Image

Whole Japanese medaka extract (30 μ g) was separated by 12% SDS-PAGE, and the membrane was blotted with PCNA antibody [HL1965] (GTX637858) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](#).

Rad1 antibody [4126]

Cat. No. GTX70317

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Applications	WB
Reactivity	Human

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000

Note : Detects a band of approximately 28 kDa

Not tested in other applications.

Calculated MW 32 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Amino terminal fragment of human Rad1 purified from E. coli.
Purification	Protein G purified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX70317 WB Image

Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with Rad1 antibody [4126] (GTX70317) diluted at 1:1000. The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



For full product information, images and publications, please visit our [website](https://www.genetex.com).