

GTX300103

Antibody panel for autophagy

Product Content

Cat No	Product Name	Reactivity	Applications	Package
GTX100685	SQSTM1 / P62 antibody [N3C1], Internal	Human, Mouse, Rat, Zebrafish, Bovine, Honeybee, Mosquito	WB, ICC/IF, IHC-P, FCM, IP, PLA	10 µl
GTX213110-01	Goat Anti-Rabbit IgG antibody (HRP)	Rabbit	WB, IHC-P, ELISA	25 µl
GTX639686	LC3B antibody [HL2798]	Human, Mouse, Rat, Zebrafish	WB, ICC/IF, IHC-P	10 µl

Note

For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 1 of 16

SQSTM1 / P62 antibody [N3C1], Internal

Cat. No. GTx100685

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, FCM, IP, PLA
Reactivity	Human, Mouse, Rat, Zebrafish, Bovine, Honeybee, Mosquito

References (105)

 Review (5)

Package

100 µl, 25 µl

PRODUCT

Summary SQSTM1 antibody, also known as ubiquitin-binding protein p62 antibody, detects the SQSTM1 protein that interacts with LC3 proteins to execute autophagosomal degradation. SQSTM1 is also reported to regulate the activation of NF κ B1 downstream signaling in addition to having other cellular functions.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
FCM	1:50-1:200
IP	1:100-1:500
PLA	Assay dependent

Not tested in other applications.

Calculated MW 48 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.39 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the center region of human SQSTM1 / P62. The exact sequence is proprietary.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 2 of 16

Purification

Purified by antigen-affinity chromatography.

Conjugation


Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES


**GTX100685 ICC/IF Image**

SQSTM1 antibody [N3C1], Internal detects SQSTM1 protein at autophagosome by immunofluorescent analysis.


Samples: HeLa cells mock (left) and treated with 50 μM Chloroquine for 24 hr (right) were fixed in 4% paraformaldehyde at RT for 15 min.

Green: SQSTM1 protein stained by SQSTM1 antibody [N3C1], Internal (GTX100685) diluted at 1:1000.

Red: Phalloidin, a F-actin marker.

**GTX100685 WB Image**

Non-transfected (-) and transfected (+) HepG2 whole cell extracts (30 μg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 antibody [N3C1], Internal (GTX100685) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


**GTX100685 WB Image**

Various whole cell extracts (30 μg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 / P62 antibody [N3C1], Internal (GTX100685) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 11 Page 3 of 16

**GTX100685 WB Image**


Untreated (–) and treated (+) NIH-3T3 whole cell extract (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 / P62 antibody [N3C1], Internal (GTX100685) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.

**GTX100685 WB Image**

Untreated (–) and treated (+) HepG2 whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 / P62 antibody [N3C1], Internal (GTX100685) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

**GTX100685 WB Image**

Untreated (–) and treated (+) Huh-7 whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 antibody [N3C1], Internal (GTX100685) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

**GTX100685 WB Image**

SQSTM1 antibody [N3C1], Internal detects SQSTM1 protein by western blot analysis.

A. 30 µg PC-12 whole cell lysate/extract

B. 30 µg Rat2 whole cell lysate/extract

10% SDS-PAGE


SQSTM1 antibody [N3C1], Internal (GTX100685) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 4 of 16

**GTX100685 WB Image**

SQSTM1 antibody [N3C1], Internal detects SQSTM1 protein by western blot analysis.

A. 30 μ g NIH-3T3 whole cell lysate/extract


B. 30 μ g JC whole cell lysate/extract

C. 30 μ g BCL-1 whole cell lysate/extract

12% SDS-PAGE

SQSTM1 antibody [N3C1], Internal (GTX100685) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


**GTX100685 IHC-P Image**

SQSTM1 / P62 antibody [N3C1], Internal detects SQSTM1 / P62 protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded human lung cancer.

SQSTM1 / P62 stained by SQSTM1 / P62 antibody [N3C1], Internal (GTX100685) diluted at 1:500.


Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

**GTX100685 IP Image**

Immunoprecipitation of SQSTM1 protein from HeLa whole cell extracts using 5 μ g of SQSTM1 antibody [N3C1], Internal (GTX100685).

Western blot analysis was performed using SQSTM1 antibody [N3C1], Internal (GTX100685).

EasyBlot anti-Rabbit IgG (GTX221666-01) was used as a secondary reagent.

**GTX100685 FCM Image**

SQSTM1 antibody [N3C1], Internal (GTX100685) detects SQSTM1 protein by flow cytometry analysis.

Sample: HeLa cell fixed in 4% paraformaldehyde at 4°C for 5 min.

Brown: Unlabelled sample was also used as a control.


Blue: SQSTM1 antibody [N3C1], Internal (GTX100685) dilution: 1:100.

Acquisition of >20,000 events were collected using Argon ion laser (488nm) and 525/30 bandpass filter.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 5 of 16



GTX100685 ICC/IF Image


SQSTM1 antibody [N3C1], Internal detects SQSTM1 protein at autophagosome by immunofluorescent analysis.

Samples: HepG2 cells treated with 3μM thapsigargin 12 hrs (right) and mock (left) were fixed in ice-cold MeOH for 10 min, permeabilize with cooled acetone for 1 min.

Green: SQSTM1 protein stained by SQSTM1 antibody [N3C1], Internal (GTX100685) diluted at 1:500.


Blue: Hoechst 33342 staining.

Scale bar = 10 μm.




GTX100685 WB Image

The data was published in the 2020 in Int J Mol Sci. [PMID: 33260618](#)



GTX100685 WB Image

The data was published in the journal Front Pharmacol in 2017. [PMID: 29326587](#)




GTX100685 WB Image

The data was published in the journal Biomolecules in 2019. [PMID: 31505885](#)




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 6 of 16




GTX100685 WB Image

The data was published in the journal Biomolecules in 2019. [PMID: 31505885](#)




GTX100685 WB Image

The data was published in the journal *Antioxidants (Basel)* in 2019. [PMID: 31906147](#)



GTX100685 WB Image

The data was published in the journal *Antioxidants (Basel)* in 2019. [PMID: 31906147](#)





GTX100685 WB Image

The data was published in the journal Sci Rep in 2017. PMID: 28134285



For full product information, images and publications, please visit our [website](#).

**GTX100685 WB Image**The data was published in the journal PLoS One in 2015.[PMID: 25946033](#)**GTX100685 WB Image**The data was published in the journal Biomolecules in 2019.[PMID: 31505885](#)

For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 8 of 16

Goat Anti-Rabbit IgG antibody (HRP)

Cat. No. GTX213110-01

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Rabbit

References (605)

Package

1 ml

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:5000-1:20000
IHC-P	1:100-1:1000
ELISA	1:5000-1:20000

Not tested in other applications.

Properties


Form	Liquid
Buffer	0.05M Tris, 0.15M NaCl
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Highly purified whole rabbit IgG
Purification	Purified by antigen-affinity chromatography.
Conjugation	Horseradish peroxidase(HRP)
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 11 Page 9 of 16

DATA IMAGES




GTX213110-01 ELISA Image

Indirect ELISA analysis performed by coating plate with recombinant SARS-CoV-2 (COVID-19) Spike S1 protein, His tag (active) (GTX135817-pro) (52.15-0.81 nM). Coated protein was probed with SARS-CoV-2 (COVID-19) Spike S1 antibody [HL134] (GTX635671) (1 µg/mL). Goat anti-rabbit IgG antibody (HRP) (GTX213110-01) (1:10000) was used to detect bound primary antibody.




GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in mouse intestine by immunohistochemical analysis. Sample: Paraffin-embedded mouse intestine. WBP11 antibody (GTX118654) diluted at 1:500. The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01) Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various amounts of denatured rabbit IgG protein were separated by 12% SDS-PAGE, and the membrane was blotted with HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) diluted at 1:5000.



GTX213110-01 IHC-P Image

WBP11 antibody detects WBP11 protein at nucleus in rat prostate by immunohistochemical analysis. Sample: Paraffin-embedded rat prostate. WBP11 antibody (GTX118654) diluted at 1:500. The signal was developed by Rabbit IgG antibody (HRP) (GTX213110-01). Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX213110-01 WB Image

Various whole cell extracts were separated by 10% SDS-PAGE, and the membrane was blotted with alpha Tubulin antibody (GTX112141) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 10 of 16




D**GTX213110-01 WB Image**The data was published in the 2022 in Cancers (Basel). [PMID: 35267540](#)

MMP1
 β -Actin

From Kim HW, et al. Cancers (Basel) (2022).
 Shown under license agreement via CiteAb.

C


From Wang F, et al. Front Aging Neurosci (2022).
 Shown under license agreement via CiteAb.

GTX213110-01 WB ImageThe data was published in the 2022 in Front Aging Neurosci. [PMID: 35264943](#)**GTX213110-01 WB Image**The data was published in the 2017 in Nat Commun. [PMID: 28281525](#)**d**

From Huang CK, et al. Nat Commun (2017).
 Shown under license agreement via CiteAb.


GTX213110-01 WB ImageThe data was published in the 2017 in Nat Commun. [PMID: 28281525](#)

For full product information, images and publications, please visit our [website](#).




GTX213110-01 WB Image

The data was published in the 2017 in *Nat Commun.* PMID: 28281525




GTX213110-01 WB Image

The data was published in the 2016 in PLoS One. [PMID: 27486856](#)




GTX213110-01 WB Image

The data was published in the journal Front Pharmacol in 2018.[PMID: 30087612](#)



GTX213110-01 WB Image

The data was published in the journal Cell Commun Signal in 2014. [PMID: 25214434](#)




GTX213110-01 WB Image


The data was published in the journal PLoS One in 2017. [PMID: 28832634](#)




For full product information, images and publications, please visit our [website](#).

**GTX213110-01 WB Image**


The data was published in the journal PLoS One in 2017. [PMID: 28945763](#)

**GTX213110-01 WB Image**


The data was published in the journal Oncoscience in 2015. [PMID: 26097885](#)

**GTX213110-01 WB Image**

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)

**GTX213110-01 WB Image**

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)

**GTX213110-01 WB Image**

The data was published in the journal EMBO Rep in 2020. [PMID: 31885126](#)



For full product information, images and publications, please visit our [website](#).

LC3B antibody [HL2798]

Cat. No. GTX639686

Host	Rabbit	Package
Clonality	Monoclonal	100 µl, 25 µl
Isotype	IgG	
Applications	WB, ICC/IF, IHC-P	
Reactivity	Human, Mouse, Rat, Zebrafish	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent

Not tested in other applications.

Observed MW (kDa) 16 (LC3B_I), 14 (LC3B_{II}, autophagosome marker) kDa.

Properties


Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide of human LC3B.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 11 Page 14 of 16

DATA IMAGES




GTX639686 WB Image

Untreated (-) and treated (+) HepG2 whole cell extracts (30 µg) were separated by 15% SDS-PAGE, and the membrane was blotted with LC3B antibody [HL2798] (GTX639686) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX639686 WB Image

Untreated (-) and treated (+) PC-12 whole cell extracts (30 µg) were separated by 15% SDS-PAGE, and the membrane was blotted with LC3B antibody [HL2798] (GTX639686) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX639686 WB Image

Untreated (-) and treated (+) NIH-3T3 whole cell extracts (30 µg) were separated by 15% SDS-PAGE, and the membrane was blotted with LC3B antibody [HL2798] (GTX639686) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX639686 IHC-P Image

LC3B antibody [HL2798] detects LC3B protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded human lung cancer.

LC3B stained by LC3B antibody [HL2798] (GTX639686) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX639686 IHC-P Image

LC3B antibody [HL2798] detects LC3B protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded rat cerebellum.


LC3B stained by LC3B antibody [HL2798] (GTX639686) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 15 of 16




GTX639686 IHC-P Image

LC3B antibody [HL2798] detects LC3B protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse cerebellum.

LC3B stained by LC3B antibody [HL2798] (GTX639686) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX639686 ICC/IF Image

LC3B antibody [HL2798] detects LC3B protein at autophagosome weak by immunofluorescent analysis.

Sample: Mock and treated HeLa cells were fixed in ice-cold MeOH for 5 min.

Green: LC3B stained by LC3B antibody [HL2798] (GTX639686) diluted at 1:1500.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.



GTX639686 IHC-P Image


LC3B antibody [HL2798] detects LC3B protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded human cerebellum.

Green: LC3B stained by LC3B antibody [HL2798] (GTX639686) diluted at 1:100.

Blue: Fluoroshield with DAPI (GTX30920).

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX639686 WB Image

Whole cell extract (30 µg) was separated by 15% SDS-PAGE, and the membrane was blotted with LC3B antibody [HL2798] (GTX639686) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody

(GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident femto Western HRP Substrate.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 16 of 16