

EpCAM antibody [ESA 214] (FITC)

Cat. No. GTX30708

| | |
|---------------------|------------------|
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Applications | ICC/IF, FCM, IHC |
| Reactivity | Human |

References (9)

Package

500 µl

Applications

Application Note

IHC, ICC (frozen or formalin-fixed paraffin-embedded (FFPE) tissue sections, cell smears. For IHC dilute conc. antibodies 1:50-1:100, use streptavidin~biotin system or polymer system, incubate 30 minutes at room temperature. IF, FITC~conjugated antibodies 10-20 µg/ml (1:10-1:20), incubate for 2 hours in the dark at RT or it can also be incubated overnight at 4°C. FITC~conjugated antibodies, 0.2-1.0 µg/0.1 ml (1:200-1:1,000)

Properties

| | |
|----------------------|----------------------------------------------------------------------------------------------------------------------------|
| Form | Liquid |
| Buffer | PBS, 1% BSA |
| Preservative | 0.05% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light. |
| Concentration | 0.20 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Small cell lung carcinoma |
| Purification | Protein A purified |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).