

Trypsin Antigen Retrieval Kit

Cat. No. GTX30934

Applications IHC-P

References (2)
Package
1 kit

PRODUCT

Summary Trypsin Antigen Retrieval Kit (2 components)

Applications

Application Note

To one ml of buffer (reagent B), add 2-4 drops of concentrated enzyme (reagent E) in a test tube, mix well. This ready-to-use reagent is good for 2-3 hours.

1. Deparaffinize FFPE tissue section is carried out as outlined in your protocol. Hydrate tissue with PBS.
2. Remove buffer, add 2-4 drops of ready-to use enzyme solution to cover the tissue section.
3. Incubate at 37°C for 5-10 minutes or 10-15 minutes at room temperature, (the time required for optimum digestion of FFPE tissue section will vary with the extent of fixation, generally approx. 5-15 minutes are sufficient at the above temperatures. In some cases it may require more time).
4. Discard trypsin reagent, wash 3-5 times with buffer.
5. The tissue sections are ready for further IHC protocol.

Properties

Form Liquid

Storage Store at 2-8°C, DO NOT FREEZE.

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).