

AEC

Cat. No. GTX30938

Applications

IHC

References (1)

Package

1 kit

PRODUCT

Summary

AEC

Applications

Application Note

To 5 ml of distilled or deionized water in a test tube, add two drop of reagent B buffer, mix well. Add one drop of reagent C chromogen, mix well. Add one drop of reagent S substrate, mix well. This ready to use reagent is good for several hours. The unused AEC solution can be discarded according to city, county, state, province or country's regulations. After the peroxidase reaction, wash slide with buffer (this buffer should not contain any sodium azide because peroxidase is inactivated), and distilled/ deionized water. Add few drops of ready to use AEC reagent, incubate at RT or 37°C for 5-10 minutes. Wash 2-5X with buffer, followed by rinsing with distilled/deionized water. Add counterstain compatible with AEC (we recommend hematoxylin counterstain for 30-60 seconds). Wash with tap water, buffer pH 7.4 or higher followed by rinsing with distilled/deionized water.

Properties

Form

Liquid

Storage

Store at 2-8°C.

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 1