

Streptavidin (FITC)

Cat. No. GTX30950

Application	ICC/IF, IHC-P, FACS	Package 1 mg
		3

PRODUCT

Summary

Streptavidin is a 55 kDa (subunit MW 14 kDa) biotin-binding protein isolated from Streptomyces avidini. Streptavidin is superior to avidin, because it does not contain carbohydrate like avidin and has no net charge at neutral pH. Streptavidin~biotin system is routinely used in Immunohistochemistry (IHC). Extinction Coefficient 1% A280=32.0. Protein content: > 95%. One mg of Streptavidin is conjugated with FITC, unconjugated Streptavidin is removed. This conjugate of streptavidin is recommended for use with Biotin~conjugated antibodies. Avidin~Biotin conjugated reagents are stable, sensitive and give less background. Fluorophore: Fluorescein isothiocyanate A max=492nm; E=520nm;

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Fluorophore/Protein: A492nm/A280nm=~1

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	Assay dependent
FACS	Assay dependent

Not tested in other applications.

PROPERTIES		
Form	Liquid	
Buffer	10mM Phosphate (pH7.4), 150mM NaCl, 10mg/ml BSA	
Preservative	0.05% Sodium azide	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.	
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)	
Conjugation	Fluorescein isothiocyanate (FITC)	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 30 Page 1 of 1