

# Interferon gamma antibody

**Cat. No. GTX31185**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, ELISA, IHC
<b>Reactivity</b>	Rat

**Package**  
100 µg

## Applications

### Application Note

We recommend the following starting dilutions: For WB: Use at a concentration of 0.1-0.2 µg/ml. For ELISA: Use at a concentration of 0.5-2.0 µg/ml. For IHC: Use at a concentration of 0.25 ug/ml. Optimal dilutions should be determined experimentally by the end user.

**Calculated MW** 18 kDa. ( [Note](#) )

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS pH7.2
<b>Preservative</b>	No preservative
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	Batch dependent (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Recombinant Rat IFN-γ
<b>Conjugation</b>	Unconjugated

### Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

## DATA IMAGES



### GTx31185 IHC Image

IHC analysis of colchicine injected rat brain (including the cortex and median eminence) tissue using IFN gamma antibody at a concentration of 0.25 mg/ml. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA) reagent.



### GTx31185 IHC Image

IHC analysis of colchicine injected rat brain (including the cortex and median eminence) tissue using IFN gamma antibody at a concentration of 0.25 mg/ml. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA) reagent.



For full product information, images and publications, please visit our [website](#).