

Bcl-2 antibody [100]

Cat. No. GTX31237

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Application	WB, IHC-P, IHC-Fr, FACS, IP
Reactivity	Human

Package
100 µg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
IHC-P	1/20-1/40
IHC-Fr	1/20-1/40
FACS	1/50-1/100
IP	Assay dependent

Note : This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Membrane permeabilisation is required for this application.

Not tested in other applications.

Calculated MW 26 kDa. ([Note](#))

PROPERTIES

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide, amino acids 41-54 of the Bcl-2 protein
Purification	Protein G purified From tissue culture supernatant



For full product information, images and publications, please visit our [website](#).


Conjugation

Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

GTx31237 WB Image

WB analysis of Jurkat cell lysate using GTx31237 Bcl-2 antibody [100].


GTx31237 IHC-P Image

IHC-P analysis of human tonsil tissue using GTx31237 Bcl-2 antibody [100].


 For full product information, images and publications, please visit our [website](#).