

HELIOS antibody

Cat. No. GTX31612

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human, Mouse, Rat

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 - 2 µg/mL
IHC-P	5 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 60 kDa. ([Note](#))

Product Note At least two isoforms are known to exist; this antibody will detect both isoforms. This antibody is predicted to not cross-react with other IKZF proteins.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	HELIOS antibody was raised against a 19 amino acid peptide near the amino terminus of human HELIOS. The immunogen is located within amino acids 50 - 100 of HELIOS.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


DATA IMAGES



GTX31612 WB Image

WB analysis of EL4 cell lysate in (A) the absence and (B) the presence of blocking peptide using GTX31612 HELIOS antibody.

Working concentration : 1 µg/ml



GTX31612 IHC-P Image

IHC-P analysis of human spleen tissue using GTX31612 HELIOS antibody.

Working concentration : 20 µg/ml



GTX31612 IHC-P Image

IHC-P analysis of human spleen tissue using GTX31612 HELIOS antibody.

Working concentration : 5 µg/ml



For full product information, images and publications, please visit our [website](https://www.genetex.com).