

Bad antibody

Cat. No. GTX31688

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, IHC-P, ELISA
Reactivity	Human, Mouse, Rat

Reference (1)

Package

100 µg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5 - 2 µg/mL
IHC-P	2 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 18 kDa. ([Note](#))

PROPERTIES

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Bad antibody was raised against a peptide corresponding to 15 amino acids near the C-terminus of human Bad.The immunogen is located within amino acids 90 - 140 of BAD.
Purification	Purified by ion exchange chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX31688 IHC-P Image

IHC-P analysis of rat thymus tissue using GTX31688 Bad antibody.
Working concentration : 10 µg/ml



GTX31688 WB Image

WB analysis of T24 cell lysate using GTX31688 Bad antibody.
Working concentration : (A) 0.5, (B) 1, and (C) 2 µg/ml



GTX31688 IHC-P Image

IHC-P analysis of rat thymus tissue using GTX31688 Bad antibody.
Working concentration : 2 µg/ml



For full product information, images and publications, please visit our [website](#).