

# Caspase 7 antibody

**Cat. No. GTX31704**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Application</b>	WB, IHC-P, ELISA
<b>Reactivity</b>	Human, Mouse, Rat

Reference ( 2 )

Package

100 µg

## APPLICATION

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5 - 1 µg/mL
IHC-P	2 µg/mL
ELISA	Assay dependent

Not tested in other applications.

**Calculated MW** 34 kDa. ( [Note](#) )

**Product Note** Depending on cell lines or tissues used, other cleavage products may be observed.

## PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	PBS
<b>Preservative</b>	0.02% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Caspase-7 antibody was raised against a 16 amino acid synthetic peptide from near the carboxy terminus of human Caspase-7. The immunogen is located within the last 50 amino acids of Caspase-7.
<b>Purification</b>	Purified by antigen-affinity chromatography
<b>Conjugation</b>	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## DATA IMAGES



**GTX31704 IHC-P Image**

IHC-P analysis of human skeletal muscle tissue using GTX31704 Caspase 7 antibody.  
Working concentration : 2 µg/ml



**GTX31704 WB Image**

WB analysis of human skeletal muscle cell lysate using GTX31704 Caspase 7 antibody.  
Working concentration : (A) 0.5, (B) 1, and (C) 2 µg/ml



For full product information, images and publications, please visit our [website](#).