

Angiopoietin 2 antibody [Angy-1-4] (Azide free)

Cat. No. GTX33914-00

Host	Human	Package
Clonality	Monoclonal	100 µg
Isotype	IgG2	
Applications	ELISA, Neutralizing/Inhibition	
Reactivity	Human	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ELISA	Assay dependent
Neutralizing/Inhibition	Assay dependent

Note : Inhibits the binding of human angiopoietin-2 to human Tie-2. ND50* = 600-800ng/ml (for 10ng/ml of Angiopoietin-2)

***ND50: 50% neutralizing dose of antibody for a given concentration of ligand (here human Angiopoietin-2).**

Not tested in other applications.

Product Note Recognizes human angiopoietin-2. Does not recognize mouse angiopoietin-2 and human angiopoietin-1.

Properties

Form	Liquid
Buffer	PBS, 10% Glycerol
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant human angiopoietin-2.
Purification	Protein A-affinity purified.
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 30 Page 1 of 2

DATA IMAGES



GTX33914-00 ELISA Image

Binding of human Angiopoietin-2 to Tie-2 (human):Fc is inhibited by the antibody anti-Angiopoietin-2 (human), mAb (rec.) (blocking) (Angy-1-4) (preservative free). Methods: Tie-2 (human):Fc was coated on an ELISA plate at 1 μ g/ml. anti-Angiopoietin-2 (human), mAb (rec.) (blocking), (Angy-1-4) (preservative free) or an unrelated mAb (recombinant) (Control) were added (starting at 40 μ g/ml with a twofold serial dilution) together with 200ng/ μ l of Angiopoietin-2 (human). After incubation for 1h at RT, the binding was detected using an anti-FLAG antibody (HRP).



GTX33914-00 Image



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 2 of 2