

Angiopoietin 2 antibody [Angy-2-1] (Azide free)

Cat. No. GTX33916-00

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Applications	ELISA, Neutralizing/Inhibition
Reactivity	Human, Mouse

References (1)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ELISA	Assay dependent
Neutralizing/Inhibition	Assay dependent

Note : Inhibits the binding of mouse angiopoietin-2 to mouse Tie-2. ND50* = 50-60ng/ml (for 10ng/ml of mouse Angiopoietin-2)

Inhibits the binding of human angiopoietin-2 to human Tie-2. ND50* = 8-12ng/ml (for 10ng/ml of human Angiopoietin-2)

***ND50: 50% neutralizing dose of antibody for a given concentration of ligand (here Angiopoietin-2).**

Not tested in other applications.

Product Note Recognizes human and mouse angiopoietin-2. Does not detect human angiopoietin-1.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant human angiopoietin-2.
Purification	Protein A-affinity purified.
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 1 of 2


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES**GTX33916-00 ELISA Image**

Binding of human Angiopoietin-2 to Tie-2 (human):Fc is inhibited by the antibody anti-Angiopoietin-2, mAb (rec.) (blocking) (Angy-2-1) (preservative free). Methods: Tie-2 (h):Fc was coated on an ELISA plate at 1 μ g/ml. anti-Angiopoietin-2, mAb (rec.) (blocking) (Angy-2-1) (preservative free) or an unrelated mAb (recombinant) (Control) were added (starting at 40 μ g/ml with a twofold serial dilution) together with 20ng/ μ l of Angiopoietin-2 (human). After incubation for 1h at RT, the binding was detected using an anti-FLAG antibody (HRP).

**GTX33916-00 Image**

For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 2 of 2