

Angiotensin 2 antibody [Angy-2-1] (Azide free)

Cat. No. GTX33916-00

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Applications	ELISA, Neutralizing /Inhibition
Reactivity	Human, Mouse

References (1)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ELISA	Assay dependent
Neutralizing /Inhibition	Assay dependent

Note : Inhibits the binding of mouse angiotensin-2 to mouse Tie-2. ND50* = 50-60ng/ml (for 10ng/ml of mouse Angiotensin-2)**Inhibits the binding of human angiotensin-2 to human Tie-2. ND50* = 8-12ng/ml (for 10ng/ml of human Angiotensin-2)*****ND50: 50% neutralizing dose of antibody for a given concentration of ligand (here Angiotensin-2).**

Not tested in other applications.

Product Note Recognizes human and mouse angiotensin-2. Does not detect human angiotensin-1.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant human angiotensin-2.
Purification	Protein A-affinity purified.
Conjugation	Unconjugated


For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX33916-00 ELISA Image

Binding of human Angiopoietin-2 to Tie-2 (human):Fc is inhibited by the antibody anti-Angiopoietin-2, mAb (rec.) (blocking) (Angy-2-1) (preservative free). Methods: Tie-2 (h):Fc was coated on an ELISA plate at 1µg/ml. anti-Angiopoietin-2, mAb (rec.) (blocking) (Angy-2-1) (preservative free) or an unrelated mAb (recombinant) (Control) were added (starting at 40µg/ml with a twofold serial dilution) together with 20ng/µl of Angiopoietin-2 (human). After incubation for 1h at RT, the binding was detected using an anti-FLAG antibody (HRP).



GTX33916-00 Image



For full product information, images and publications, please visit our [website](#).