

MYH9 antibody [SF9]

Cat. No. GTX33939

Host	Human
Clonality	Monoclonal
Isotype	IgG2
Applications	WB, ICC/IF, ELISA, EM
Reactivity	Human, Mouse, Rat, Drosophila

References (1)
Package
100 µg

Applications

Application Note

Immunocytochemistry: 1:1000, Western blot: 1:1000

Product Note Recognizes human, mouse, rat and drosophila myosin IIA (heavy chain).

Properties

Form	Liquid
Buffer	PBS, 10% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length myosin IIA from rat liver.
Purification	Protein A-affinity purified.
Purity	≥95% by SDS-PAGE.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX33939 Image



GTX33939 ICC/IF Image

Human Myosin IIA (non-muscle) (heavy chain) is detected by immunocytochemistry using anti-myosin IIA (non-muscle) (rec.) (SF9). Method: HeLa cells are grown in standard culture conditions, fixed with methanol, and incubated with anti-Myosin IIA (non-muscle) (rec.) (SF9) (1 μ g/ml in PBS-BSA). After incubation for 30 min at RT and several washes in PBS, cells are treated with a goat anti-human (Cy3) antibody for 30 min at RT, washed and mounted in Moewiol. Nuclei are stained with DAPI.



For full product information, images and publications, please visit our [website](#).