Survivin antibody [Mix]

Cat. No. GTX34059

Host	Mouse
Clonality	Multiclonal
lsotype	lgG1
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:2000
IHC-P	1:200-1:500
Not tested in other applications.	

Package 100 μl

Calculated MW

16 kDa. (<u>Note</u>)

Properties		
Form	Liquid	
Buffer	PBS, 0.5% BSA, 50% Glycerol	
Preservative	0.02% Sodium azide	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)	
Immunogen	Recombinant Protein	
Purification	Purified by antigen-affinity chromatography	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	



For full product information, images and publications, please visit our <u>website</u>.

DATA IMAGES



GTX34059 IHC-P Image

IHC-P analysis of human tonsil tissue using GTX34059 Survivin antibody [Mix]. Negative control (the lower left coner) was secondary antibody only.


Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min) Dilution : 1:200



GTX34059 IHC-P Image

IHC-P analysis of mouse spleen tissue using GTX34059 Survivin antibody [Mix]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min) Dilution : 1:200



GTX34059 IHC-P Image

IHC-P analysis of human stomach cancer tissue using GTX34059 Survivin antibody [Mix]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min) Dilution : 1:200



GTX34059 IHC-P Image

IHC-P analysis of rat spleen tissue using GTX34059 Survivin antibody [Mix]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min) Dilution : 1:200



GTX34059 IHC-P Image

IHC-P analysis of rat brain tissue using GTX34059 Survivin antibody [Mix]. Negative control (the lower left coner) was secondary antibody only. Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)

Dilution : 1:200



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 09 / 16 Page 2 of 2