

IKB beta antibody [1F3]

Cat. No. GTX34098

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Applications	IHC-P
Reactivity	Human, Rat

Package 100 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100-1:200

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS, 0.5% BSA, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant Protein
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Conjugation	Unconjugated For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 1 of 2

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com



DATA IMAGES




GTX34098 IHC-P Image

IHC-P analysis of human stomach tissue using GTX34098 IKB beta antibody [1F3]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval: Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)


Dilution: 1:200



GTX34098 IHC-P Image

IHC-P analysis of human kidney tissue using GTX34098 IKB beta antibody [1F3].

Dilution: 1:200




GTX34098 IHC-P Image

IHC-P analysis of human liver cancer tissue using GTX34098 IKB beta antibody [1F3]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval: Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)

Dilution: 1:200



GTX34098 IHC-P Image

IHC-P analysis of human lung carcinoma tissue using GTX34098 IKB beta antibody [1F3].

Dilution: 1:200



GTX34098 IHC-P Image

IHC-P analysis of rat spleen tissue using GTX34098 IKB beta antibody [1F3]. Negative control (the lower left coner) was secondary antibody only.

Antigen retrieval: Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)

Dilution: 1:200



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 2 of 2