

Caspase 9 antibody [3-20]

Cat. No. GTX34168

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, IHC-P, IP
Reactivity	Human, Mouse, Rat

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:5000
IHC-P	Assay dependent
IP	1:200

Not tested in other applications.

Calculated MW 46 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 0.5% BSA, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic Peptide
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 1 of 2


DATA IMAGES

**GTX34168 IHC-P Image**

IHC-P analysis of mouse kidney tissue using GTX34168 Caspase 9 antibody [3-20]. Negative control (the lower left corner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)


Dilution : 1:200

**GTX34168 IHC-P Image**

IHC-P analysis of rat kidney tissue using GTX34168 Caspase 9 antibody [3-20]. Negative control (the lower left corner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)


Dilution : 1:200

**GTX34168 IHC-P Image**

IHC-P analysis of human stomach cancer tissue using GTX34168 Caspase 9 antibody [3-20]. Negative control (the lower left corner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)


Dilution : 1:200

**GTX34168 IHC-P Image**

IHC-P analysis of rat spleen tissue using GTX34168 Caspase 9 antibody [3-20]. Negative control (the lower left corner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)

Dilution : 1:200

**GTX34168 IHC-P Image**

IHC-P analysis of rat lung tissue using GTX34168 Caspase 9 antibody [3-20]. Negative control (the lower left corner) was secondary antibody only.

Antigen retrieval : Sodium citrate pH6.0 was used for antibody retrieval (>98°C, 20min)

Dilution : 1:200



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 2 of 2