

c-Kit antibody [C117/370]

Cat. No. GTX34606

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Applications	IHC-P
Reactivity	Human

Package $100 \, \mu g$

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1-2μg/ml for 30 minutes at RT

Note: Staining of formalin-fixed tissues require heating tissue sections in 1mM EDTA buffer, pH 9.5, for 45 min at 95°C followed by cooling at RT for 20 minutes.

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS, 0.05% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant human CD117 protein
Purification	Protein A/G purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 12 / 06 Page 1 of 2




DATA IMAGES



GTX34606 IHC-P Image

IHC-P analysis of human gastrointestinal stromal tumor (GIST) tissue using GTX34606 c-Kit antibody [C117/370].



GTX34606 IHC-P Image

IHC-P analysis of human gastrointestinal stromal tumor (GIST) tissue using GTX34606 c-Kit antibody [C117/370].



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 06 Page 2 of 2