

## EpCAM antibody [EGP40/1110]

## Cat. No. GTX34691

Host	Mouse	Package
Clonality	Monoclonal	100 µg
Isotype	IgG2b	
Applications	IHC-P, Protein Array	
Reactivity	Human, Mouse, Rat	

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1-2µg/ml for 30 min at RT
Protein Array	Assay dependent

**Note : Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes.**

Not tested in other applications.

## Properties


Form	Liquid
Buffer	PBS, 0.05% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A recombinant fragment from the cytoplasmic domain (around aa 280-350) of human EpCAM (exact sequence is proprietary)
Purification	Protein A/G purified
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 28 Page 1 of 2


## DATA IMAGES

**GTX34691 IHC-P Image**


IHC-P analysis of rat colon tissue using GTX34691 EpCAM antibody [EGP40/1110].

**GTX34691 IHC-P Image**

IHC-P analysis of human ovarian carcinoma tissue using GTX34691 EpCAM antibody [EGP40/1110].

**GTX34691 Protein Array Image**

Analysis of Protein Array containing more than 19,000 full-length human proteins using EpCAM Mouse Monoclonal Antibody (EGP40/1110). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.

**GTX34691 IHC-P Image**

IHC-P analysis of rat lung tissue using GTX34691 EpCAM antibody [EGP40/1110].



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 28 Page 2 of 2