

# Moesin antibody [MSN/492]

## Cat. No. GTX34848

Host	Mouse	
Clonality	Monoclonal	
Isotype	lgG1	
Applications	WB, ICC/IF, IHC-P, FCM	
Reactivity	Human	

Package 100 μg

# Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1-2μg/ml
ICC/IF	1-2μg/ml
IHC-P	1-2μg/ml for 30 minutes at RT
FCM	1-2μg/10 <sup>6</sup> cells

Note: Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes.

Not tested in other applications.

Calculated MW 68 kDa. (Note)

**Product Note** We do not recommend use of this product for Rat samples.

Properties	
Form	Liquid
Buffer	PBS, 0.05% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant full-length human Moesin protein
Purification	Protein A/G purified
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 13 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## **DATA IMAGES**




## GTX34848 IHC-P Image

IHC-P analysis of human testicular carcinoma tissue using GTX34848 Moesin antibody [MSN/492].



## GTX34848 IHC-P Image

IHC-P analysis of human melanoma tissue using GTX34848 Moesin antibody [MSN/492].



## GTX34848 WB Image

WB analysis of Jurkat cell lysate using GTX34848 Moesin antibody [MSN/492].



For full product information, images and publications, please visit our website.

Date 2025 / 05 / 13 Page 2 of 2