

CaMKII (oxidized) antibody

Cat. No. GTX36254

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr, PLA
Reactivity	Human, Mouse, Rat

References (31)

Package

50 µl

Applications

Application Note

For WB, use at a dilution of 1:1,000, For IF, use at a dilution of 1:50 with paraffin-embedded sections and 1:100 with frozen sections.

Product Note

Mouse. Predicted to react with human based on 100% sequence identity.

Properties

Form	Liquid
Buffer	Serum
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	CaMKII synthetic peptide with oxidized Met281/282
Purification	Unpurified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX36254 IHC-P Image

Immunofluorescent staining of heart sections from mice treated with saline, AngII, or Isoproterenol and probed for oxidized or total CaMKII. Red staining is positive for oxidized or total CaMKII, and blue staining is positive for nuclei. Mice lacking a critical subunit of NADPH oxidase (p47^{-/-}) did not show increased levels of oxidized CaMKII in response to AngII.




GTX36254 IHC-P Image

Immunofluorescent staining of heart sections from WT and MsrA^{-/-} mice (a model of heart disease) treated with AngII and probed for oxidized (GTX36254) or total CaMKII and blue staining for nuclei.




GTX36254 WB Image

Immunoblot of oxidized WT CaMKII probed with antiserum against oxidized M281/282 with increasing ratios of oxidized peptide.



GTX36254 WB Image

Immunoblot of WT CaMKII and M281/282V mutant after no treatment, oxidation, or autophosphorylation probed with antibodies against total, autophosphorylated, or oxidized M281/282 CaMKII (GTX36254).



GTX36254 WB Image

WB analysis of various sample lysates using using GTX36254 CaMKII (oxidized) antibody.

- Lane 1 : Recombinant CaMKII
 - Lane 2 : Untreated mouse heart cell lysates
 - Lane 3 : H2O2-treated mouse heart cell lysates
- Dilution : 1:1000



For full product information, images and publications, please visit our [website](http://www.genetex.com).