

CDC2 (phospho Thr161) antibody

Cat. No. GTX38597

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

References (1)

Package

50 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:1000
IHC-P	1:50-1:100

Not tested in other applications.

Calculated MW	34 kDa. (Note)
Product Note	CDC2 (phospho-Thr161) antibody detects endogenous levels of CDC2 only when phosphorylated at threonine 161.

Properties

Form	Liquid
Buffer	PBS, 150mM NaCl, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	The antiserum was produced against synthesized phosphopeptide derived from human CDC2 around the phosphorylation site of threonine161 (T -Y-T(p)-H-E).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX38597 WB Image

WB analysis of COLO205 cell (Lane 1 and 2) and K562 cell (Lane 3) lysates using GTX38597 CDC2 (phospho Thr161) antibody.



GTX38597 IHC-P Image

IHC-P analysis of human breast carcinoma tissue using GTX38597 CDC2 (phospho Thr161) antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).