

Swc8 antibody [MIL3]

Cat. No. GTX39351

Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Applications	IHC-Fr, FCM
Reactivity	Pig

Package 2 ml

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	1/10-1/50
FCM	1/20

Note : Use $50\mu l$ of the suggested working dilution to label 10^6 cells in $100\mu l$.

Not tested in other applications.

Properties	
Form	Liquid
Buffer	Tissue culture supernatant, 200mM Tris-HCl, 5-10% FBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Immunogen	Porcine Lamina Propria Leucocytes
Purification	Tissue Culture Supernatant
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 12 Page 1 of 1