

MICA/MICB antibody [6D4] (Low endotoxin, azide free)

Cat. No. GTX39659

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Applications	IHC-Fr, FCM, IP, Neutralizing /Inhibition
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	1/100
FCM	Assay dependent
IP	Assay dependent
Neutralizing /Inhibition	Assay dependent

Note : Clone 6D4 is reported to inhibit the cytotoxicity of NK cells stimulated by IFN alpha-treated dendritic cells (Jinushi et al. 2003).

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	MICA transfected C1R cells.
Purification	Protein G purified From tissue culture supernatant
Endotoxin	< 0.01 EU/µg (determined by the LAL assay)
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).