

CDC2 antibody

Cat. No. GTX40151

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB
Reactivity	Xenopus laevis

References (1)
Package
100 µg

Applications

Application Note

WB: Use at an assay dependent dilution. Predicted molecular weight: 34 kDa. Optimal dilutions/concentrations should be determined by the end user.

Calculated MW	35 kDa. (Note)
Product Note	3E1.1 reacts strongly with immunoblotted antigen but does not immunoprecipitate native p34cdc2.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	C terminal fragment, starting from methionine 85 of Xenopus Cdc2 .
Purification	IgG fraction
Conjugation	Unconjugated

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).