

Neisseria meningitidis antibody (FITC)

Cat. No. GTX40370

| | |
|---------------------|------------------------|
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | ICC/IF |
| Reactivity | Neisseria meningitidis |

Package
1 ml

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| ICC/IF | 1:10-1:50 |

Note : Acetone fixation of the antigen source is recommended prior to staining.

Not tested in other applications.

Properties

| | |
|----------------------|--|
| Form | Liquid |
| Buffer | PBS, 1% BSA |
| Preservative | 0.1% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light. |
| Concentration | 4-5 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Whole cell prep. of N. meningitidis types A, B & C |
| Purification | IgG fraction |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).