

CD86 antibody [24F]

Cat. No. GTX41790

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-Fr, FCM, IP
Reactivity	Rat

Package
250 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	Assay dependent
FCM	1/25-1/200
IP	Assay dependent

Note : Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl.

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	HTLV-1 transformed Lewis-S1 cells.
Purification	Protein G purified From tissue culture supernatant
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX41790 IHC-Fr Image

IHC-Fr analysis of rat lymph node tissue using GTX41790 CD86 antibody [24F].




GTX41790 IHC-Fr Image

IHC-Fr analysis of rat lymph node tissue using GTX41790 CD86 antibody [24F].



GTX41790 FCM Image

FACS analysis of rat splenocytes using GTX41790 CD86 antibody [24F].



GTX41790 IHC-Fr Image

IHC-Fr analysis of rat lymph node tissue using GTX41790 CD86 antibody [24F].



For full product information, images and publications, please visit our [website](#).