

CD8 antibody [YTS169.4]

Cat. No. GTX41817

Host	Rat
Clonality	Monoclonal
Isotype	IgG2b
Applications	IHC-Fr, FCM
Reactivity	Mouse

Package

25 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	Assay dependent
FCM	1/50-1/100

Note : Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl.

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Purification	Protein G purified From tissue culture supernatant
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX41817 IHC-Fr Image


IHC-Fr analysis of mouse lymph node tissue using GTX41817 CD8 antibody [YTS169.4].


GTX41817 IHC-Fr Image

IHC-Fr analysis of mouse lymph node tissue using GTX41817 CD8 antibody [YTS169.4].


GTX41817 IHC-Fr Image

IHC-Fr analysis of mouse lymph node tissue using GTX41817 CD8 antibody [YTS169.4].


GTX41817 FCM Image

FACS analysis of mouse spleen cells using GTX41817 CD8 antibody [YTS169.4].



For full product information, images and publications, please visit our [website](#).