

CD5L antibody, C-term

Cat. No. GTX41904

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5-1µg/ml
IHC-P	Assay dependent

Note : This antibody detects a band of approximately 38kDa in Raji cell lysates.

Not tested in other applications.

Calculated MW	38 kDa. (Note)
Product Note	This antibody detects an epitope within the C-terminal region of human CD5L.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	13 amino acid peptide sequence near the carboxy terminus of human CD5L.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).


Date 2026 / 10 / 06 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX41904 IHC-P Image

IHC-P analysis of human lymph node tissue using GTX41904 CD5L antibody, C-term.



GTX41904 WB Image

WB analysis of Raji cell lysate using GTX41904 CD5L antibody at (A)0.5, (B)1 and (C)2 ug/ml.



For full product information, images and publications, please visit our [website](#).