

## Coagulation factor III/Tissue Factor antibody [TF9-10H10]

## Cat. No. GTX42410

Host	Mouse	Package
Clonality	Monoclonal	100 µg
Isotype	IgG1	
Applications	WB, ICC/IF, IHC-Fr, FCM	
Reactivity	Human	

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-Fr	Assay dependent
FCM	Assay dependent

Not tested in other applications.

Calculated MW 33 kDa. ([Note](#))

Properties	
<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 200mM Mannitol
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Denatured Tissue factor isolated from human brain by the Factor VII affinity method (Guha et al. 1986).
<b>Purification</b>	Protein G purified
<b>Conjugation</b>	Unconjugated
<b>Note</b>	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 2



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 2 of 2