

## ICAM2 antibody [CBRIC2/2] (Low endotoxin, azide free)

**Cat. No. GTX42520**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2a
<b>Applications</b>	IHC-Fr, FCM, IP, Functional Assay
<b>Reactivity</b>	Human

References ( 1 )

Package

100 µg

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	Assay dependent
FCM	1/100
IP	Assay dependent
Functional Assay	Assay dependent

**Note : Use 10µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl.**

Not tested in other applications.

**Product Note**

This antibody recognizes human ICAM2. Clone CBRIC2/2 is reported to inhibit interactions between ICAM2 and LFA-1 (De Fougerolles et al. 1991).

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Transfected COS cells expressing ICAM-2 cDNA.
<b>Purification</b>	Protein G purified From tissue culture supernatant
<b>Endotoxin</b>	< 0.01 EU/µg (determined by the LAL assay)
<b>Conjugation</b>	Unconjugated



For full product information, images and publications, please visit our [website](#).

Date 2026 / 05 / 04 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).