

## Bmi1 antibody

## Cat. No. GTX45791

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, ELISA, Multiplexing
Reactivity	Human

Package  
50 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:13000
ICC/IF	1:200
ELISA	1:5000-1:30000
Multiplexing	Assay dependent

Not tested in other applications.

Calculated MW 37 kDa. ( [Note](#) )

## Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.91 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide corresponding to amino acids 252-264 of human Bmi1 protein.
Purification	Purified by antigen-affinity chromatography. From serum
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 2


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

**DATA IMAGES****GTX45791 WB Image**

Western blot using GeneTex's Affinity Purified anti-Bmi1 antibody shows detection of a band ~37 kDa corresponding to human Bmi1 (arrowhead). Approximately 20 µg of a U2OS whole cell lysate was separated by 4-20% SDS-PAGE and transferred onto nitrocellulose. After blocking in PBS containing 5% nonfat dry milk, the membrane was probed overnight at 4° C with the primary antibody diluted to 1:1,000 in PBS containing 1% nonfat dry milk. The membrane was washed and reacted with a 1:20,000 dilution of IRDye800 conjugated rabbit anti-Goat IgG [H&L] for 45 min at room temperature. IRDye800 fluorescence image was captured using the OdysseyR Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.

**GTX45791 ICC/IF Image**


Immunofluorescence using affinity purified goat anti Bmi1(GTX45791) shows nuclear staining (green) of methanol fixed (100%, 5 min) HepG2 cells. The cells were blocked and permeabilized in 1%BSA / 10% normal donkey serum / 0.3M glycine in 0.1% PBS-Tween for 1h prior to incubation with the primary antibody (1:200 dilution) overnight at +4°C and detected with a 488nm fluorescent dye conjugated secondary Ab. Cell nuclei are stained with DAPI (blue) and plasma membranes are stained with WGA (red).

**GTX45791 WB Image**

WB analysis of U2OS whole cell lysate using GTX45791 Bmi1 antibody.

Loading : 20 µg

Dilution : 1:1000

**GTX45791 ICC/IF Image**

ICC/IF analysis of MeOH-fixed HepG2 cells using GTX45791 Bmi1 antibody.

Green : Primary antibody

Red : Wheat germ agglutinin

Blue : DAPI

Dilution : 1:200



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 2 of 2