

TSPAN3 antibody, Internal

Cat. No. GTx46420

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 µg/ml
IHC-P	2-10 µg/ml

Not tested in other applications.

Calculated MW 28 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to an Internal region of Human TSPAN3
Purification	Affinity Purified
Conjugation	Unconjugated
	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 1 of 2

DATA IMAGES

**GTX46420 WB Image**

WB analysis of HeLa cells using GTX46420 TSPAN3 antibody at 0.2-1 μ g/ml.

**GTX46420 IHC-P Image**

IHC-P analysis of human testis tissue using GTX46420 TSPAN3 antibody at 1:600. Left to right : DAPI, TSPAN3, Merge. Heat-induced antigen retrieval method utilizing Sodium Citrate buffer.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 2 of 2