

TSPAN5 antibody, Internal

Cat. No. GTx46421

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 µg/ml
IHC-P	2-10 µg/ml

Not tested in other applications.

Calculated MW 30 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to an Internal region of Human TSPAN5
Purification	Affinity Purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 10 Page 1 of 2

DATA IMAGES




GTX46421 WB Image

WB analysis of Jurkat cells using GTX46421 TSPAN5 antibody at 0.25 μ g/ml.

Lane A : marker

Lane B : Jurkat cells



GTX46421 IHC-P Image

IHC-P analysis of human kidney tissue using GTX46421 TSPAN5 antibody at 4.0-8.0 μ g/ml.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 10 Page 2 of 2