

## NOLC1 antibody, C-term

Cat. No. GTX46486

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, IHC-P, IHC-Wm
<b>Reactivity</b>	Human, Zebrafish

Package  
100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 ug/ml
IHC-P	2-10 ug/ml
IHC-Wm	Assay dependent

Not tested in other applications.

**Calculated MW** 74 kDa. ([Note](#))

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 2% Sucrose
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	A synthetic peptide corresponding to a C-terminal region of Human NOLC1
<b>Purification</b>	Protein A purified
<b>Conjugation</b>	Unconjugated

## Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


## DATA IMAGES

**GTX46486 WB Image**

WB analysis of HepG2 cells using GTX46486 NOLC1 antibody at 2.5 $\mu$ g/ml.

Lane A : marker

Lane B : HepG2 cells

**GTX46486 IHC-P Image**

IHC-P analysis of human kidney tissue using GTX46486 NOLC1 antibody at 4.0-8.0 $\mu$ g/ml.

**GTX46486 IHC-Wm Image**

IHC-Wm analysis of zebrafish embryo (27 hpf) using GTX46486 NOLC1 antibody at 1:250.



For full product information, images and publications, please visit our [website](#).