

SLC33A1 antibody, Internal

Cat. No. GTX46828

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 ug/ml
IHC-P	2-10 ug/ml

Not tested in other applications.

Calculated MW 61 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to an Internal region of Human SLC33A1
Purification	Affinity Purified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX46828 WB Image

WB analysis of human B lymphoblast cells using GTX46828 SLC33A1 antibody at 1µg/ml.




GTX46828 WB Image

WB analysis of human placenta tissue using GTX46828 SLC33A1 antibody at 0.2-1µg/ml.



GTX46828 IHC-P Image

IHC-P analysis of small intestine tissue using GTX46828 SLC33A1 antibody at 5.0µg/ml.



GTX46828 WB Image

WB analysis of HeLa cells using GTX46828 SLC33A1 antibody at 1µg/ml.



For full product information, images and publications, please visit our [website](#).